arXiv:math/0405383v2 [math.RT] 1 Jun 2004

REPRESENTATIONS OF THE RATIONAL CHEREDNIK ALGEBRAS OF DIHEDRAL TYPE

TATYANA CHMUTOVA

Abstract. In this paper we describe the Jordan-Hölder series of the standard modules over the rational Cherednik algebras associated with the dihedral group. In particular, we compute the characters of the irreducible representations from the category $\mathcal{O}$ and classify the finite dimensional representations.

1. Introduction

The rational Cherednik algebra $H_{\mathsf{k}}(W)$ for a complex reflection group W arises from the study of Dunkl operators ([Du],[DO]), and is an important special case of symplectic reflection algebras ([EG]).

In the recent years, a number of authors studied representations of $H_{\mathsf{k}}(W)$ (see e.g. [GGOR, BEG, BEG1, DO, De, Go]). Specifically, the paper [GGOR] defines the category $\mathcal{O}$ for $H_{\mathsf{k}}(W)$, and shows that its basic structure is quite similar to the structure of the category $\mathcal{O}$ for a simple Lie algebra, studied by Bernstein, I. Gelfand and S. Gelfand. In particular, the main problem about this category is to find the characters of the irreducible modules in $\mathcal{O}$ and their multiplicities in standard modules.

Although many partial results are available, this problem remains open in all interesting situations. The goal of this paper is to solve this problem for dihedral groups, which is the simplest nontrivial case. We describe the structure of the standard modules over rational Cherednik algebras corresponding to the dihedral group $I_2(d)$. The structure depends on the parity of d.

When d is odd the Cherednik algebra depends on one complex parameter. For general value of the parameter, all the standard modules are irreducible. The set of exceptional values consists of a finite number of arithmetic progressions of rational numbers, each with the common difference 1. For these values, the standard modules have length at most 2, and at least one of them is reducible. The category $\mathcal{O}$ is not semisimple in this case. This classification is given in theorems 3.1.1 and 3.1.2.

The case of even d is more complicated since the Cherednik algebra depends on two complex parameters. If the values of the parameters do not belong to four exceptional families of lines, all the standard modules are irreducible. For generic points of the exceptional lines the maximal length of standard modules is 2 and at least one is reducible. The category $\mathcal{O}$ for these values of parameters is not semisimple. At the points of intersection of the lines in these families, the structure of the standard modules might be complicated, and the length might reach 4. For some of the intersection points even the category of the finite dimensional modules is not semisimple. This case is described in theorems 3.2.2 and 3.2.3.

2. Preliminaries

2.1. Let W be a finite real reflection group, $\mathfrak{h}_{\mathbb{R}}$ be its reflection representation, and $\mathfrak{h}$ be the complexification of this representation. Fix a non-degenerate W-invariant scalar product $(-,-)$ on $\mathfrak{h}$. Denote by the same sign the induced scalar product on $\mathfrak{h}^*$. Let $S \subset W$ be the set of all reflections $s \in W$. To each reflection $s \in S$ we associate its fixed hyperplane H_s, a nonzero linear function $\alpha_s \in \mathfrak{h}^*$ which vanishes on H_s, and the element $\alpha_s^\vee = \frac{2(\alpha_s,-)}{(\alpha_s,\alpha_s)} \in \mathfrak{h}$. In other words, to each s we associate the corresponding positive root α_s and coroot $\alpha_s^\vee$.

Let us fix a W-invariant function $\mathsf{k} : S \to \mathbb{C}$, where W acts on S via conjugation.

Definition 2.1.1. *The rational Cherednik algebra* $H_{\mathsf{k}}(W)$ is an associative algebra generated by all elements of W, $\mathfrak{h}^*$, and $\mathfrak{h}$ with the following defining relations:

$$wxw^{-1} = w(x), \quad wyw^{-1} = w(y), \quad [x,x'] = 0, \quad [y,y'] = 0,$$

$$[y,x] = (y,x) - \sum_{s\in S} \mathsf{k}(s)(y,\alpha_s)(\alpha_s^\vee, x)s,$$

where $x, x' \in \mathfrak{h}^*$, $y, y' \in \mathfrak{h}$ and $w \in W$ (see also [EG]).

2.2. For rational Cherednik algebras there exists an analog of Verma modules (see [BEG]), called *standard* modules. They are indexed by irreducible representations of W. Let V be an irreducible representation of W. It can be considered as a representation of $\mathbb{C}[W] \ltimes \mathbb{C}[\mathfrak{h}^*]$, where $y \in \mathfrak{h}$ acts by 0.

Definition 2.2.1. *The standard module* $M_{\mathsf{k}}(V)$ over $H_{\mathsf{k}}(W)$ corresponding to V is defined by the formula

$$M_{\mathsf{k}}(V) = H_{\mathsf{k}}(W) \otimes_{\mathbb{C}[W]\ltimes\mathbb{C}[\mathfrak{h}^*]} V.$$

Definition 2.2.2. The standard module $M_{\mathsf{k}}(V)$ and its quotients are called *modules with lowest W-weight V.*

As a vector space, $M_{\mathsf{k}}(V)$ is naturally identified with $\mathbb{C}[\mathfrak{h}] \otimes V$. Hence the usual grading on $\mathbb{C}[\mathfrak{h}]$ induces a grading on $M_{\mathsf{k}}(V)$.

The standard module $M_{\mathsf{k}}(V)$ has a unique simple quotient $L_{\mathsf{k}}(V)$ (see [DO]).

An important special case of standard modules is $M_{\mathsf{k}}(\mathsf{triv})$, which is also called the *polynomial representation* of $H_{\mathsf{k}}(W)$. The space of this representation is $\mathbb{C}[\mathfrak{h}]$. An element $w \in W$ acts on $\mathbb{C}[\mathfrak{h}]$ in the natural way, $x \in \mathfrak{h}$ acts by left multiplication, and the action of $y \in \mathfrak{h}$ is given by the *Dunkl operator*

$$D_y = \partial_y - \sum_{s\in S} \mathsf{k}(s)\frac{(y,\alpha_s)}{\alpha_s}(1-s). \tag{1}$$

Definition 2.2.3. A vector in $M_{\mathsf{k}}(V)$ is called *singular* if it is annihilated by every $y \in \mathfrak{h}$.

In particular, the singular vectors in $M_{\mathsf{k}}(\mathsf{triv})$ are exactly those annihilated by all Dunkl operators.

2.3. Every rational Cherednik algebra contains a canonically defined $\mathfrak{sl}_2$-triple:

$$\mathbf{h} = \frac{1}{2}\sum_i (x_i y_i + y_i x_i), \quad E = \frac{1}{2}\sum_i x_i^2, \quad F = -\frac{1}{2}\sum_i y_i^2,$$

where $\{x_i\}$ and $\{y_i\}$ are dual orthonormal bases of $\mathfrak{h}^*$ and $\mathfrak{h}$. Therefore, every lowest weight $H_{\mathsf{k}}(W)$-module M is also an $\mathfrak{sl}_2$-module from the category $\mathcal{O}$. The element $\mathbf{h}$ is W-invariant and satisfies the relations

$$[\mathbf{h}, x] = x, \quad [\mathbf{h}, y] = -y$$

for all $x \in \mathfrak{h}^*$ and $y \in \mathfrak{h}$. The spectrum of $\mathbf{h}$ on M is $\mathbb{Z}_{\geqslant 0} + h_0$, where h_0 is the lowest weight of $\mathbf{h}$ on M.

Remark 2.3.1. The generalized eigenspace of $\mathbf{h}$ with eigenvalue $h_0 + i$ consists of all the elements of degree i in M.

As in the case of Lie groups, one can define the category $\mathcal{O}$ of $H_{\mathsf{k}}(W)$-modules.

Definition 2.3.2. *The category* $\mathcal{O}$ is the category of $H_{\mathsf{k}}(W)$-modules M, such that M is a direct sums of finite dimensional generalized eigenspaces of $\mathbf{h}$, and the spectrum of $\mathbf{h}$ is bounded from below.

Standard modules $M_{\mathsf{k}}(V)$ and their quotients are objects of the category $\mathcal{O}$. It is known that $\{L_{\mathsf{k}}(V)\}_{V \in Irrep(W)}$ are precisely the simple objects of $\mathcal{O}$.

For $M \in \mathcal{O}$, let $M = \oplus_a M_a$ be its $\mathbf{h}$-weight decomposition.

Definition 2.3.3. *The character* of M is a formal power series in t given by

$$\chi_M(g,t) = \sum_a \mathrm{Tr}\big|_{M_a}(g) t^a,$$

where $g \in W$, and $\mathrm{Tr}\big|_{M_a}(g)$ is the trace of g-action on M_a.

The character of the standard module $M_{\mathsf{k}}(V)$ is given by

$$\chi_{M_{\mathsf{k}}(V)}(g,t) = \frac{\chi_V(g) t^{h_0(V)}}{\det\big|_{\mathfrak{h}}(1-gt)},$$

where $h_0(V) = \frac{1}{2}\dim\mathfrak{h} - \left(\sum_{s\in S} \mathsf{k}(s)s\right)\big|_V$ is the lowest weight of $\mathbf{h}$ on $M_{\mathsf{k}}(V)$ (see [BEG], formula (1.5)).

Remark 2.3.4. For any character $\varepsilon : W \to \mathbb{C}^*$ there is a symmetry between $H_{\mathsf{k}}(W)$ and $H_{\varepsilon\mathsf{k}}(W)$. Namely, there exists an isomorphism $H_{\mathsf{k}}(W) \xrightarrow{\sim} H_{\varepsilon\mathsf{k}}(W)$, which acts identically on $\mathfrak{h}$ and $\mathfrak{h}^*$, and sends $w \in W$ to $\varepsilon(w)w$. This isomorphism induces an equivalence of categories of representations of these algebras. Under this equivalence, $M_{\mathsf{k}}(V)$ goes to $M_{\varepsilon\mathsf{k}}(\varepsilon \otimes V)$.

3. Results

The group $I_2(d)$ acts via reflections on $\mathfrak{h}_{\mathbb{R}} \simeq \mathbb{C}$. The subset $S \subset W$ of reflections consists of elements $s_j : z \mapsto \omega^j \bar{z}$, where $\omega = e^{2\pi i/d}$ and $1 \leqslant j \leqslant d$.

3.1. **The case $d = 2m+1$.** All the reflections in S are conjugate, and an $I_2(m)$-invariant function $\mathsf{k} : S \to \mathbb{C}$ is a constant function $\mathsf{k} = c \in \mathbb{C}$. Denote by H_c the corresponding rational Cherednik algebra $H_{\mathsf{k}}(W)$.

The irreducible representations of $I_2(d)$ are: the trivial representation triv, the sign representation sgn, and two-dimensional representations τ_l, where $1 \leqslant l \leqslant m$. The representations τ_l are realized on the span of $\{z^l, \bar{z}^l\}$.

In what follows, we denote the product dc by r.

Theorem 3.1.1. *The structure of $M_c(\mathsf{triv})$ for different values of c is described in the table below.*

Values of c	**Structure of the standard module**	**Character of the irreducible quotient**
$0 < r \equiv \pm l \pmod d$, *where* $1 \leqslant l \leqslant m$	*$M_c(\mathsf{triv})$ has a submodule isomorphic to $L_c(\tau_l)$. The quotient is irreducible of dimension r^2 .*	$t^{1-r}\dfrac{\det\big\|_{\tau_l}(1-gt^r)}{\det\big\|_{\mathfrak{h}}(1-gt)}$
$0 < c = n + \frac{1}{2}$, *where* $n \in \mathbb{Z}$	*$M_c(\mathsf{triv})$ has a submodule isomorphic to $M_c(\mathsf{sgn})$. The quotient is irreducible and infinite dimensional.*	$t^{1-r}\dfrac{1-\mathsf{sgn}(g)t^{2r}}{\det\big\|_{\mathfrak{h}}(1-gt)}$
Other values of c	*$M_c(\mathsf{triv})$ is irreducible.*	$t^{1-r}\dfrac{1}{\det\big\|_{\mathfrak{h}}(1-gt)}$

The description of the $M_c(\mathsf{sgn})$ follows from theorem 3.1.1 by symmetry (Remark 2.3.4).

Theorem 3.1.2. *Fix $1 \leqslant l \leqslant m$. The structure of $M_c(\tau_l)$ for different values of c is described in the following table.*

Values of c	**Structure of the standard module**	**Character of the irreducible quotient**
$0 < r \equiv \pm l \pmod d$	*$M_c(\tau_l)$ has a submodule isomorphic to $M_c(\mathsf{sgn})$. The quotient is irreducible and infinite dimensional.*	$t\dfrac{\chi_{\tau_l}(g)-t^r}{\det\big\|_{\mathfrak{h}}(1-gt)}\mathsf{sgn}(g)$
$0 > r \equiv \pm l \pmod d$	*$M_c(\tau_l)$ has a submodule isomorphic to $M_c(\mathsf{triv})$. The quotient is irreducible and infinite dimensional.*	$t\dfrac{\chi_{\tau_l}(g)-t^{-r}}{\det\big\|_{\mathfrak{h}}(1-gt)}$
Other values of c	*$M_c(\tau_l)$ is irreducible.*	$t\dfrac{\chi_{\tau_j}(g)}{\det\big\|_{\mathfrak{h}}(1-gt)}$

Remark 3.1.3. All the standard modules over H_c are of length either 1 or 2.

3.2. **The case $d = 2m$.** There are two conjugacy classes of reflections: $\{s_{2i}\}_{1\leqslant i\leqslant m}$ and $\{s_{2i-1}\}_{1\leqslant i\leqslant m}$. Hence a W-invariant function $\mathsf{k} : S \to \mathbb{C}$ is a pair (k_1, k_2), where $k_1 = \mathsf{k}(s_{2i+1})$ and $k_2 = \mathsf{k}(s_{2i})$. Let H_{k_1,k_2} denote the corresponding rational Cherednik algebra $H_{\mathsf{k}}(W)$.

The group $I_2(d)$ has four irreducible one-dimensional representations: triv, sgn, ε_1, $\varepsilon_2 = \varepsilon_1 \otimes \mathsf{sgn}$, and $(m-1)$ two-dimensional representations τ_l, $1 \leqslant l < m$. The representation ε_1 is defined by

$$\varepsilon_1(s_{2i+1}) = 1, \quad \varepsilon_1(s_{2i}) = -1,$$

and τ_l's are realized on $\mathrm{span}\{z^l, \bar{z}^l\}$.

Remark 3.2.1. Note that $\mathsf{sgn} \otimes \tau_l = \tau_l$ and $\varepsilon_i \otimes \tau_l = \tau_{m-l}$ for $i = 1, 2$.

The structure of the standard H_{k_1,k_2}-modules depends on the location of the point (k_1, k_2) with respect to certain exceptional lines on the complex plane of parameters (k_1, k_2). We distinguish the four families E_r^+, E_r^-, L_i^1, L_i^2 of parallel exceptional lines:

- for every integer r, non divisible by m, let

$$E_r^+ := \{(k_1, k_2) : k_1 + k_2 = r/m\}, \qquad E_r^- := \{(k_1, k_2) : k_1 - k_2 = r/m\} \ ;$$

- for every integer i, let

$$L_i^1 := \{(k_1, k_2) : k_1 = i + 1/2\}, \qquad L_i^2 := \{(k_1, k_2) : k_2 = i + 1/2\} \ .$$

By symmetry reasons, in order to describe the standard modules induced from one-dimensional representations, it is enough to describe $M_{k_1,k_2}(\mathsf{triv})$.

Theorem 3.2.2. *Depending on the location of the point (k_1, k_2), there are following seven cases for the standard module $M_{k_1,k_2}(\mathsf{triv})$.*

(i) $(k_1, k_2) \in E_r^+$ **for some** $r > 0$**, and** $(k_1, k_2) \notin L_i^1, L_i^2$ **for all** $i > 0$**.**

In this case determine an integer l in the range $1 \leqslant l < m$ from the congruence $r \equiv \pm l \pmod{2m}$. Then $M_{k_1,k_2}(\mathsf{triv})$ contains a submodule isomorphic to $L_{k_1,k_2}(\tau_l)$. The quotient is irreducible and has the character

$$\chi_{L_{k_1,k_2}(\mathsf{triv})}(g, t) \quad = \quad t^{1-r} \frac{\det\big|_{\tau_l}(1 - g t^r)}{\det\big|_{\mathfrak{h}}(1 - gt)} \ .$$

In particular, its dimension is r^2.

(ii) $(k_1, k_2) \in L_i^1$ **for some** $i > 0$**, and** $(k_1, k_2) \notin E_r^+, L_{i'}^2$ **for all** $r, i' > 0$**.**

In this case the module $M_{k_1,k_2}(\mathsf{triv})$ contains a submodule isomorphic to $L_{k_1,k_2}(\varepsilon_2) \simeq M_{k_1,k_2}(\varepsilon_2)$. The quotient is irreducible and infinite dimensional. The character is given by the formula

$$\chi_{L_{k_1,k_2}(\mathsf{triv})}(g, t) \quad = \quad t^{1-m(k_1+k_2)} \frac{1 - t^{2mk_1}\chi_{\varepsilon_2}(g)}{\det\big|_{\mathfrak{h}}(1 - gt)} \ .$$

(iii) $(k_1,k_2)\in L^2_{i'}$ **for** $i'>0$**, and** $(k_1,k_2)\notin E^+_r, L^1_i$ **for all** $r,i>0$**.**

In this case the module $M_{k_1,k_2}(\mathsf{triv})$ *contains a submodule isomorphic to* $L_{k_1,k_2}(\varepsilon_1)\simeq M_{k_1,k_2}(\varepsilon_1)$*. The quotient is irreducible and infinite dimensional. The character is given by the formula*

$$\chi_{L_{k_1,k_2}(\mathsf{triv})}(g,t) \quad = \quad t^{1-m(k_1+k_2)}\frac{1-t^{2mk_2}\chi_{\varepsilon_1}(g)}{\det\big|_{\mathfrak{h}}(1-gt)}\ .$$

(iv) $(k_1,k_2)\in E^+_r\cap L^1_i$ **(resp.** $(k_1,k_2)\in E^+_r\cap L^2_{i'}$**) for some** $r,i,i'>0$**, and** $k_1<k_2$ **(resp.** $k_2<k_1$**).**

Determine an integer l *from the same conditions* $r\equiv\pm l \pmod{2m}$ *and* $1\leqslant l<m$*. We have the following inclusions of the submodules*

$$M_{k_1,k_2}(\mathsf{triv})\supset M_{k_1,k_2}(\varepsilon_2)\supset L_{k_1,k_2}(\tau_l)$$

$$(\textit{resp.}\quad M_{k_1,k_2}(\mathsf{triv})\supset M_{k_1,k_2}(\varepsilon_1)\supset L_{k_1,k_2}(\tau_l)\ )\ .$$

The quotient $M_{k_1,k_2}(\varepsilon_2)/L_{k_1,k_2}(\tau_l)$ *(resp.* $M_{k_1,k_2}(\varepsilon_1)/L_{k_1,k_2}(\tau_l)$*), is irreducible. The quotient* $Q_{k_1,k_2}=M_{k_1,k_2}(\mathsf{triv})/M_{k_1,k_2}(\varepsilon_2)$ *(resp.* $Q_{k_1,k_2}=M_{k_1,k_2}(\mathsf{triv})/M_{k_1,k_2}(\varepsilon_1)$*) contains a submodule* P_{k_1,k_2} *isomorphic to* $L_{k_1,k_2}(\varepsilon_1)$ *(resp.* $L_{k_1,k_2}(\varepsilon_2)$*). However, the double quotient* $Q_{k_1,k_2}/P_{k_1,k_2}$ *is irreducible, i.e. isomorphic to* $L_{k_1,k_2}(\mathsf{triv})$*.*

In both cases the character of $L_{k_1,k_2}(\mathsf{triv})$ *is given by the formula*

$$\chi_{L_{k_1,k_2}(\mathsf{triv})}(g,t) \quad = \quad t^{1-r}\frac{1+t^{2r}\mathsf{sgn}(g)-t^{2mk_1}\varepsilon_2(g)-t^{2mk_2}\varepsilon_1(g)}{\det\big|_{\mathfrak{h}}(1-gt)}\ .$$

In particular, its dimension is equal to $4m^2k_1k_2$*.*

(v) $(k_1,k_2)\in E^+_r\cap L^1_i$ **(resp.** $(k_1,k_2)\in E^+_r\cap L^2_{i'}$**) for some** $r,i,i'>0$**, and** $k_1>k_2$ **(resp.** $k_2>k_1$**).**

Again, let l *be an integer, satisfying the conditions* $r\equiv\pm l \pmod{2m}$ *and* $1\leqslant l<m$*. Then* $M_{k_1,k_2}(\mathsf{triv})$ *contains a submodule isomorphic to* $L_{k_1,k_2}(\varepsilon_2)$ *(resp.* $L_{k_1,k_2}(\varepsilon_1)$*). The quotient* $Q_{k_1,k_2}=M_{k_1,k_2}(\mathsf{triv})/L_{k_1,k_2}(\varepsilon_2)$ *(resp.* $Q_{k_1,k_2}=M_{k_1,k_2}(\mathsf{triv})/L_{k_1,k_2}(\varepsilon_1)$*) contains a submodule* P_{k_1,k_2} *isomorphic to* $L_{k_1,k_2}(\tau_l)$*. The quotient* $Q_{k_1,k_2}/P_{k_1,k_2}$ *is irreducible and has the character*

$$\chi_{L_{k_1,k_2}(\mathsf{triv})}(g,t) \quad = \quad t^{1-r}\frac{\det\big|_{\tau_l}(1-gt^r)}{\det\big|_{\mathfrak{h}}(1-gt)}\ .$$

In particular, its dimension is r^2*.*

(vi) $(k_1,k_2)\in L^1_i\cap L^2_{i'}$ **for some** $i,i'>0$**.**

In this case the module $M_{k_1,k_2}(\mathsf{triv})$ *contains a submodule isomorphic to* $L_{k_1,k_2}(\mathsf{sgn})\simeq M_{k_1,k_2}(\mathsf{sgn})$*. The quotient* $Q_{k_1,k_2}=M_{k_1,k_2}(\mathsf{triv})/L_{k_1,k_2}(\mathsf{sgn})$ *contains two submodules* $P^1_{k_1,k_2}\simeq L_{k_1,k_2}(\varepsilon_1)$ *and* $P^2_{k_1,k_2}\simeq L_{k_1,k_2}(\varepsilon_2)$*, such that* $P^1_{k_1,k_2}\cap P^2_{k_1,k_2}=0$*. The quotient* $Q_{k_1,k_2}/(P^1_{k_1,k_2}\oplus P^2_{k_1,k_2})$ *is irreducible, i.e. isomorphic to* $L_{k_1,k_2}(\mathsf{triv})$ *with the character*

$$\chi_{L_{k_1,k_2}(\mathsf{triv})}(g,t) \quad = \quad t^{1-r}\frac{1+t^{2r}\mathsf{sgn}(g)-t^{2mk_1}\varepsilon_2(g)-t^{2mk_2}\varepsilon_1(g)}{\det\big|_{\mathfrak{h}}(1-gt)}\ .$$

In particular, it has dimension $4m^2k_1k_2$*.*

(vii) For all other values of (k_1, k_2)
the module $M_{k_1,k_2}(\mathsf{triv})$ *is irreducible and has the character*

$$\chi_{L_{k_1,k_2}(\mathsf{triv})}(g,t) \quad = \quad t^{1-r}\frac{1}{\det\big|_{\mathfrak{h}}(1-gt)} \ .$$

The structure of the standard modules, induced from the two-dimensional representations τ_l of the group $W = I_2(d)$ is described by the following theorem.

Theorem 3.2.3. *Fix* $1 \leqslant l < m$ *and consider the two-dimensional representation* τ_l *of* $I_2(d)$*. Then, depending on the location of the point* (k_1, k_2)*, there are following four cases for the standard module* $M_{k_1,k_2}(\tau_l)$*.*

(i) $(k_1, k_2) \in E_r^+$ **for some** $r \equiv \pm l \pmod{2m}$**, and** $(k_1, k_2) \notin E_{r'}^-$ **for all** $r' \equiv m \pm l \pmod{2m}$**.**

The module $M_{k_1,k_2}(\tau_l)$ *contains a submodule* I_{k_1,k_2} *generated by one vector of degree* $|r|$*.*

If $r > 0$*, then* $I_{k_1,k_2} \simeq M_{k_1,k_2}(\mathsf{sgn})$*. If* $r < 0$*, then* $I_{k_1,k_2} \simeq M_{k_1,k_2}(\mathsf{triv})$*.*

The quotient $M_{k_1,k_2}(\tau_l)/I_{k_1,k_2}$ *is irreducible and infinite dimensional. If* $r > 0$*, its character is equal to*

$$\chi_{L_{k_1,k_2}(\tau_l)}(g,t) \quad = \quad t\frac{\chi_{\tau_l}(g) - t^r}{\det\big|_{\mathfrak{h}}(1-gt)}\mathsf{sgn}(g) \ .$$

For $r < 0$*, the character can be easily obtained using symmetry arguments.*

(ii) $(k_1, k_2) \notin E_r$ **for all** $r \equiv \pm l \pmod{2m}$**, but** $(k_1, k_2) \in E_{r'}^-$ **for some** $r' \equiv m \pm l \pmod{2m}$**.**

The module $M_{k_1,k_2}(\tau_l)$ *contains a submodule* $\widetilde{I}_{k_1,k_2}$ *generated by one vector of degree* $|r'|$*.*

If $r' > 0$*, then* $\widetilde{I}_{k_1,k_2} \simeq M_{k_1,k_2}(\varepsilon_2)$*. If* $r' < 0$*, then* $\widetilde{I}_{k_1,k_2} \simeq M_{k_1,k_2}(\varepsilon_1)$*.*

The quotient $M_{k_1,k_2}(\tau_l)/\widetilde{I}_{k_1,k_2}$ *is irreducible. If* $r' > 0$*, its character is equal to*

$$\chi_{L_{k_1,k_2}(\tau_l)}(g,t) \quad = \quad t\frac{\chi_{\tau_l}(g) - t^{r'}}{\det\big|_{\mathfrak{h}}(1-gt)}\varepsilon_2(g) \ .$$

For $r' < 0$*, the character can be easily obtained using symmetry arguments.*

(iii) $(k_1, k_2) \in E_r^+ \cap E_{r'}^-$**, for some** $r \equiv \pm l \pmod{2m}$ **and** $r' \equiv m \pm l \pmod{2m}$**.**

The module $M_{k_1,k_2}(\tau_l)$ *contains two submodules:* I_{k_1,k_2} *and* $\widetilde{I}_{k_1,k_2}$*.*

If $\min\{|k_1|, |k_2|\} \in \mathbb{Z} + 1/2$*, then one of these submodules contains another one; otherwise they intersect by* 0*. The quotient of* $M_{k_1,k_2}(\tau_l)$ *by those two submodules is irreducible in both cases. If* $r, r' > 0$*, then the character is*

$$\chi_{L_{k_1,k_2}(\tau_l)}(g,t) \quad = \quad t\frac{\chi_{\tau_l}(g) - t^r}{\det\big|_{\mathfrak{h}}(1-gt)}\mathsf{sgn}(g)$$

in the first case, and

$$\chi_{L_{k_1,k_2}(\tau_l)}(g,t) \quad = \quad t\frac{\chi_{\tau_l}(g) - t^r\mathsf{sgn}(g) - t^{r'}\varepsilon_2(g)}{\det\big|_{\mathfrak{h}}(1-gt)}$$

in the second case.
For other combinations of signs of r and r', the characters can be easily obtained using symmetry arguments.

(iv) For all other values of (k_1, k_2)
the module $M_{k_1,k_2}(\tau_l)$ is irreducible and has the character

$$\chi_{L_{k_1,k_2}(\tau_l)}(g,t) \quad = \quad t\frac{\tau_l(g)}{\det\big|_{\mathfrak{h}}(1-gt)} \ .$$

4. Examples.

In order to illustrate the Theorems 3.2.2 and 3.2.3, let us look at $W = I_2(8)$. The Theorem 3.2.2 gives us the following picture for $M_{k_1,k_2}(\mathsf{triv})$ in (k_1,k_2)-plane.



Notations:

— – $M_{k_1,k_2}(\mathsf{triv}) \supset L_{k_1,k_2}(\tau_i)$ for some i, the quotient is finite dimensional;

- - - – $M_{k_1,k_2}(\mathsf{triv}) \supset M_{k_1,k_2}(\varepsilon_1)$, quotient is generically infinite dimensional;

⋯⋯ – $M_{k_1,k_2}(\mathsf{triv}) \supset M_{k_1,k_2}(\varepsilon_2)$, quotient is generically infinite dimensional;

▪ – $M_{k_1,k_2}(\mathsf{triv})$ contains $M_{k_1,k_2}(\varepsilon_1)$ and $M_{k_1,k_2}(\varepsilon_2)$, quotient is finite dimensional.

For all the points outside the indicated lines, $M_{k_1,k_2}(\mathsf{triv})$ is irreducible.

Using this information and the symmetry arguments, one can find all the values of (k_1,k_2), for which there exist finite dimensional representation of H_{k_1,k_2}:



The following table lists the finite dimensional irreducible representations for $k_1, k_2 > 0$.

Values of parameters		**Finite dimensional irreducible modules**	**Dimension**
(k_1,k_2) is generic point on E_r^+		$L_{k_1,k_2}(\mathsf{triv})$	r^2
(k_1,k_2) is generic point on $E_{r'}^-$	$r' > 0$	$L_{k_1,k_2}(\varepsilon_1)$	${r'}^2$
	$r' < 0$	$L_{k_1,k_2}(\varepsilon_2)$	
$(k_1,k_2) \in E_r^+ \cap E_{r'}^-$, denoted by $\odot$ on the picture above	$r' > 0$	$L_{k_1,k_2}(\mathsf{triv})$, $L_{k_1,k_1}(\varepsilon_1)$	$r^2 - {r'}^2$, ${r'}^2$
	$r' < 0$	$L_{k_1,k_2}(\mathsf{triv})$, $L_{k_1,k_1}(\varepsilon_2)$	
All the other $(k_1,k_2) \in E_r^+ \cap E_{r'}^-$	$r' > 0$	$L_{k_1,k_2}(\mathsf{triv})$, $L_{k_1,k_1}(\varepsilon_1)$	r^2, ${r'}^2$
	$r' < 0$	$L_{k_1,k_2}(\mathsf{triv})$, $L_{k_1,k_1}(\varepsilon_2)$	
Isolated points on the picture above		$L_{k_1,k_2}(\mathsf{triv})$	$64k_1k_2$

Remark 4.0.4. If (k_1, k_2) lies on one of the exceptional lines, the category $\mathcal{O}$ is not semisimple.

If (k_1, k_2) is one of the points denoted by $\odot$ in the picture above, the category of finite dimensional H_{k_1,k_2}-modules is not semisimple. For example, in the case of $k_1 > k_2 > 0$ $\quad \mathrm{Ext}^1(L_{k_1,k_2}(\mathsf{triv}), L_{k_1,k_2}(\varepsilon_1)) \neq 0$.

Applying the theorem 3.2.3 to $M_{k_1,k_2}(\tau_1)$, we will get the following picture:



Notations:

—— – $M_{k_1,k_2}(\tau_1) \supset I_{k_1,k_2}$ and $I_{k_1,k_2} \simeq M_{k_1,k_2}(\mathsf{sgn})$;

- - - - – $M_{k_1,k_2}(\tau_1) \supset I_{k_1,k_2}$ and $I_{k_1,k_2} \simeq M_{k_1,k_2}(\mathsf{triv})$;

-·-·-·- – $M_{k_1,k_2}(\tau_1) \supset \widetilde{I}_{k_1,k_2}$ and $\widetilde{I}_{k_1,k_2} \simeq M_{k_1,k_2}(\varepsilon_2)$;

·········· – $M_{k_1,k_2}(\tau_1) \supset \widetilde{I}_{k_1,k_2}$ and $\widetilde{I}_{k_1,k_2} \simeq M_{k_1,k_2}(\varepsilon_1)$;

▪ – $I_{k_1,k_2} \cap \widetilde{I}_{k_1,k_2} = 0$;

$\odot$ – $I_{k_1,k_2} \subset \widetilde{I}_{k_1,k_2}$;

⬡ – $I_{k_1,k_2} \supset \widetilde{I}_{k_1,k_2}$.

As before, for all the points outside the indicated lines, $M_{k_1,k_2}(\tau_1)$ is irreducible.

5. Preparations for the proofs.

In this section we collect known results, which we need for the proofs.

5.1. **The Knizhnik-Zamolodchikov functor.** We will briefly remind the construction of the KZ-functor which, maps certain modules over the Cherednik algebra to modules over the Artin braid group. A more detailed discussion on this subject can be found in [BEG1], [GGOR] and [BMR].

5.1.1. *Hecke algebras and the Knizhnik-Zamolodchikov functor.* Let us first recall the definition of Hecke algebras associated to a reflection group W.

Let $\mathfrak{h}_{reg} = \mathfrak{h} \setminus \bigcup_{s\in S} H_s$. The W-action on $\mathfrak{h}$ preserves $\mathfrak{h}_{reg}$. Fix $x_0 \in \mathfrak{h}_{reg}$.

Definition 5.1.1. The *Artin braid group* associated to W is $B_W = \pi_1(\mathfrak{h}_{reg}/W, x_0)$.

One can describe B_W using generators and relations as follows (see [Br]). Fix a chamber C in $\mathfrak{h}$, and let Σ be the set of reflections with respect to the walls of this chamber. The generators T_s are numbered by element $s \in \Sigma$. An element T_s corresponds to the monodromy around the wall of C, fixed by s. The relations are

$$\underbrace{T_sT_tT_s\dots}_{m(s,t)\text{ factors}} = \underbrace{T_tT_sT_t\dots}_{m(s,t)\text{ factors}} \quad \text{for all } s,t \in S,$$

where $m(s,t)$ is the order of the element st.

Let $\{q_s\}_{s\in\Sigma}$ be the set of parameters such that $q_s = q_t$ if s and t are conjugated to each other.

Definition 5.1.2. The *Hecke algebra* $\mathcal{H}(\{q_s\})$ of a reflection group W with parameters q_s is the quotient of the group algebra of the Artin braid group B_W by the relations

$$(T_s - 1)(T_s + q_s) = 0.$$

Now let us define the Knizhnik-Zamolodchikov functor KZ. For any H_{k}-module M, finitely generated over $\mathbb{C}[\mathfrak{h}]$, define its localization M_{loc} by

$$M_{loc} = M \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}_{reg}].$$

This module has a natural action of $H_{\mathsf{k}\ loc}$.

The polynomial representation (see section 2.2) gives an injective homomorphism $H_{\mathsf{k}} \hookrightarrow W \ltimes \mathcal{D}(\mathfrak{h}_{reg})$, which extends to an isomorphism $H_{\mathsf{k}\ loc} \xrightarrow{\sim} W \ltimes \mathcal{D}(\mathfrak{h}_{reg})$. Hence the module M_{loc} has a structure of $W \ltimes \mathcal{D}(\mathfrak{h}_{reg})$-module, and therefore, a structure of W-equivariant $\mathcal{D}$-module on $\mathfrak{h}_{reg}$.

Since M is finitely generated over $\mathbb{C}[\mathfrak{h}]$, the module M_{loc} is a vector bundle over $\mathfrak{h}_{reg}$. This vector bundle is equipped with a connection ∇, coming from the $\mathcal{D}$-module structure. The horizontal sections of this connection define a monodromy representation

$$\rho_M : \pi_1(\mathfrak{h}_{reg}/W, x_0) \to GL(\mathbb{C}^N),$$

where N is the rank of M_{loc}.

Definition 5.1.3. A functor $\mathsf{KZ} : M \mapsto \rho_M$ is called *the Knizhnik-Zamolodchikov functor.*

$$\mathsf{KZ} : \left(\begin{array}{c} H_{\mathsf{k}}(W)\text{-modules finitely generated} \\ \text{over } \mathbb{C}[\mathfrak{h}] \end{array}\right) \to \left(\begin{array}{c} \text{Finite dimensional} \\ B_W\text{-modules} \end{array}\right)$$

In case of dihedral groups the images of the standard modules under the action of KZ-functor were studied in details by C. Dunkl in [Du1].

5.1.2. *Properties of the* KZ*-functor.* Let M be an H_{k}-module. If $q_s = e^{-2\pi i\mathsf{k}(s)}$ for all s, then the action of B_W on $\mathsf{KZ}(M)$ factors through the action of $\mathcal{H}(\{q_s\})$. If M is the standard module corresponding to a representation V of the group W, the obtained representation of $\mathcal{H}(\{q_s\})$ is the one corresponding to V via Tits' Deformation Theorem.

The following facts about the KZ-functor are important for us.

Proposition 5.1.4. [GGOR] *The functor* KZ *is exact.*

Theorem 5.1.5. ([GGOR], Theorem 5.13) *The functor* $\mathsf{KZ} : \mathcal{O} \to B_W$*-mod factors through a functor* $\mathsf{KZ} : \mathcal{O}/\mathcal{O}_{tor} \to \mathcal{H}(\{q_s\})$*-mod, where* $\mathcal{O}_{tor}$ *is the category of modules, supported on the union of reflection hyperplanes.*

Theorem 5.1.6. ([BEG1] Lemma 2.10) *Let* N *be an* H_{k}*-module, which is torsion free over* $\mathbb{C}[\mathfrak{h}]$*. Then for any* $M \in \mathcal{O}$ *the canonical map*

$$Hom_{H_{\mathsf{k}}}(M, N) \to Hom_{\mathcal{H}}(\mathsf{KZ}(M), \mathsf{KZ}(N))$$

is injective.

5.2. **Singular vectors in the polynomial representation.** Let M be a module from the category $\mathcal{O}$. Denote by U the space of singular vectors in M. This space is nonzero, since the action of y's decreases the eigenvalues of $\mathbf{h}$ by 1, and the spectrum of $\mathbf{h}$ is bounded from below. From the defining relations of Cherednik algebra follows that U is preserved under the W-action.

Proposition 5.2.1. *A module* $M \in \mathcal{O}$ *is irreducible if and only if* U *spans* M *and is an irreducible representation of* W*.*

□

Hence, in order to describe the structure of standard modules, it is important to study U.

Denote by $U' \subset U$ the space of all singular vectors of positive degree. For the polynomial representation, this space was studied by C. Dunkl, E. Opdam and M.F.E. de Jeu.

Theorem 5.2.2. (see [DJO]) *Consider the Cherednik algebra* $H_c(I_2(d))$*, where* $d = 2m + 1$*.*

1. *Let* $c = n \pm l/d > 0$*, where* n *and* l *are integers and* $1 \leqslant l \leqslant m$*. Then* $U' \subset M_c(\mathsf{triv})$ *is isomorphic to* τ_l *as a representation of* W*. This space sits in degree* $dn + l$*. It is spanned by*

$$f = \oint \left(\frac{1}{w}\right)^{n+\frac{2l}{d}} z^l (w - \bar{z}^d)^{n+\frac{l}{d}} (w - z^d)^{n+\frac{l}{d}-1} dw$$

   and its conjugate.
2. *Let* $c = n + 1/2 > 0$*. Then* $U' \subset M_c(\mathsf{triv})$ *is isomorphic to the sign representation of* W*, and is spanned by the singular vector* $g = (z^d - \bar{z}^d)^{2n+1}$*.*
3. *For all other values of* c*, there are no singular vectors of positive degree in* $M_c(\mathsf{triv})$*.*

Theorem 5.2.3. (see [DJO]) *Consider the Cherednik algebra* $H_{k_1,k_2}(I_2(2m))$*. The space* U' *is nonzero only in the following cases.*

(1) *If* $(k_1, k_2) \in E_r^+$ *for some* $0 < r \equiv \pm l \pmod{2m}$ *where* $1 \leqslant l < m$, *then* $U' \subset M_{k_1,k_2}(\mathsf{triv})$ *contains a* W*-subrepresentation isomorphic to* τ_l *and consisting of vectors of degree* r.

(2) *If* $(k_1, k_2) \in L_i^1$ *for some* $i \geqslant 0$, *then* $U' \subset M_{k_1,k_2}(\mathsf{triv})$ *contains a* W*-subrepresentation, isomorphic to* ε_2 *and spanned by the singular vector* $(z^m - \bar{z}^m)^{2i+1}$.

(3) *If* $(k_1, k_2) \in L_{i'}^2$ *for some* $i' \geqslant 0$, *then* $U' \subset M_{k_1,k_2}(\mathsf{triv})$ *contains a* W*-subrepresentation isomorphic to* ε_1 *and spanned by the singular vector* $(z^m + \bar{z}^m)^{2i+1}$.

(4) *If* $(k_1, k_2) \in L_i^1 \cap L_{i'}^2$ *for some* $i, i' \geqslant 0$, *then* $U' \subset M_{k_1,k_2}(\mathsf{triv})$ *is isomorphic to* $\varepsilon_1 \oplus \varepsilon_2 \oplus \mathsf{sgn}$ *as a* W*-representation. It is spanned by singular vectors* $(z^m + \bar{z}^m)^{2i+1}$, $(z^m - \bar{z}^m)^{2i+1}$, *and* $(z^m - \bar{z}^m)^{2i+1}(z^m + \bar{z}^m)^{2i'+1}$.

Note that if (k_1, k_2) *belongs to the intersection of* E_r^+ *with some of the lines* L_i^1 *(resp.* L_i^2 *), then* U' *is the direct sum of two irreducible representations of* W, *which are described in the cases* (1) *and* (2) *(resp.* (3)*). The intersection of lines* L_i^1 *and* $L_{i'}^2$ *is described separately in the case* (4), *because an additional singular vector appears.*

5.3. **The lowest h-weights of standard modules.** Let $V \subset U \subset M$ be an irreducible W-subrepresentation, consisting of singular vectors. Then V is contained in the generalized eigenspace of $\mathbf{h}$ with eigenvalue $h_0(V)$. Indeed, the ideal generated by V is an $H_{\mathsf{k}}(W)$-module of lowest weight V. Hence, the smallest eigenvalue of $\mathbf{h}$ on this module is $h_0(V)$, and it is achieved on the generating vectors, i.e. on V.

The knowledge of the spectrum of $\mathbf{h}$ will help us to understand U as a representation of W.

Proposition 5.3.1. (1) *Let* $d = 2m + 1$. *The lowest weights of standard* H_c*-modules are given in the table below:*

$M_c(\mathsf{triv})$	$M_c(\mathsf{sgn})$	$M_c(\tau_l)$
$1 - dc$	$1 + dc$	1

(2) *Let* $d = 2m$. *The lowest weights of standard* H_{k_1,k_2}*-modules are given in the table below:*

$M_{k_1,k_2}(\mathsf{triv})$	$M_{k_1,k_2}(\varepsilon_1)$	$M_{k_1,k_2}(\mathsf{sgn})$	$M_{k_1,k_2}(\varepsilon_2)$	$M_{k_1,k_2}(\tau_l)$
$1 - R$	$1 - R'$	$1 + R$	$1 + R'$	1

where $R = m(k_1 + k_2)$ *and* $R' = m(k_1 - k_2)$.

Proof. The proposition follows from the formula for $h_0(V)$ (see section 2.3). □

5.4. **Some irreducible finite dimensional representations.**

Proposition 5.4.1. ([CE], Theorem 2.3) *Let* W *be a real reflection group, and* $V \subset M_{\mathsf{k}} = M_{\mathsf{k}}(\mathsf{triv})$ *be a* W*-subrepresentation of dimension* $d = \dim(\mathfrak{h})$ *sitting in degree* r *and consisting of singular vectors. Let* I_{k} *be the ideal generated by* V. *Assume that the quotient representation* $Q_{\mathsf{k}} = M_{\mathsf{k}}/I_{\mathsf{k}}$ *is finite dimensional. Then* *the representation* Q_{k} *is irreducible. Its character is given by the formula*

$$\chi_{Q_{\mathsf{k}}}(g,t) = t^{h_0(\mathsf{triv})} \frac{\det\big|_V (1-gt^r)}{\det|_{\mathfrak{h}^*}(1-gt)}.$$

In particular, the dimension of Q_{k} *is* r^d.

6. Proofs

Remark 6.0.2. In all the classification theorem the character formulas follow easily from the structure of standard modules.

6.1. Case of $d = 2m+1$.

Proof of theorem 3.1.2. Let us describe the W-action on the set U' of singular vectors of positive degree.

Claim 1. *There are no two-dimensional irreducible subrepresentations in* U'.

By proposition 5.3.1, $\mathbf{h}$ acts by 1 on all irreducible two-dimensional representations of W consisting of singular vectors. However, the space on which $\mathbf{h}$ acts by 1, is exactly the space of vectors of degree 0, which does not intersect U'.

Claim 2. *For* $c \neq n \pm l/d$ *there are no one-dimensional representations in* U'.

By symmetry reasons (see remark 2.3.4), it is enough to prove that U' does not contain the sign representation.

Suppose U' contains a W-representation isomorphic to sgn. This representation generates an H_c-submodule isomorphic to $M_c(\mathsf{sgn})$ in $M_c(\tau_l)$. Therefore

$$\mathrm{Hom}_{H_c}(M_c(\mathsf{sgn}), M_c(\tau_l)) \neq 0.$$

Both $M_c(\mathsf{sgn})$ and $M_c(\tau_l)$ are free over $\mathbb{C}[\mathfrak{h}]$. By theorem 5.1.6,

$$\dim \mathrm{Hom}_{H_{\mathsf{k}}}(M_c(\mathsf{sgn}), M_c(\tau_l)) \leqslant \dim \mathrm{Hom}_{\mathcal{H}}(\mathsf{KZ}(M_c(\mathsf{sgn})), \mathsf{KZ}(M_c(\tau_l))), \tag{2}$$

and hence,

$$\mathrm{Hom}_{\mathcal{H}}(\mathsf{KZ}(M_c(\mathsf{sgn})), \mathsf{KZ}(M_c(\tau_l))) \neq 0. \tag{3}$$

Let us compute the dimension of $\mathrm{Hom}_{\mathcal{H}\text{-mod}}(\mathsf{KZ}(M_c(\mathsf{sgn})), \mathsf{KZ}(M_c(\tau_l)))$.

Recall that in case of $W = I_2(d)$ the Hecke algebra $\mathcal{H}(q)$ is generated by two elements T_1 and T_2 with relations

$$(T_j - 1)(T_j + q) = 0, \quad \underbrace{T_1 T_2 \dots T_1}_{d \text{ factors}} = \underbrace{T_2 T_1 \dots T_2}_{d \text{ factors}}.$$

In order for theorems 5.1.5 and 5.1.6 to hold, we need $q = e^{-2\pi i c}$.

Representation $\mathsf{KZ}(M_c(\tau_l))$ of $\mathcal{H}(q)$ corresponds to τ_l via Tits' Deformation Theorem. If $0 < c \neq n \pm l/d$, this representation is given explicitly by:

$$T_1 \mapsto \begin{pmatrix} 1 & 0 \\ c_l & -q \end{pmatrix}; \quad T_2 \mapsto \begin{pmatrix} -q & c_l' \\ 0 & 1 \end{pmatrix},$$

where $c_l c_l' = q(2 + \omega^l + \omega^{-l})$ (see [GP], p. 268; note that in our case the sign of the generators is different). Representation $\mathsf{KZ}(M_c(\mathsf{sgn}))$ is given explicitly by $T_1 \mapsto -q$ and $T_2 \mapsto -q$.

Hence, $\dim\big[\mathrm{Hom}_{\mathcal{H}}(\mathsf{KZ}(M_c(\mathsf{sgn})), \mathsf{KZ}(M_c(\tau_l)))\big] = 0$ for $0 < c \neq n \pm l/d$. This contradicts to (3), and hence U' does not contain subrepresentations isomorphic to sgn. Claim 2 is proved.

From Claims 1 and 2 it follows that $U' = 0$ for $r \not\equiv \pm l \pmod d$. Thus, U is the space of all vectors of degree 0 in $M_c(\tau_l)$. It is isomorphic to τ_l as a representation of W, and spans $M_c(\tau_l)$. By proposition 5.2.1, $M_c(\tau_l)$ is irreducible for these values of c. Hence, the last row of the table is proved.

Claim 3. *If* $0 < c = n \pm l/d$ *for some* $n \in \mathbb{Z}$, *then* $U' \simeq \mathsf{sgn}$ *as a* W*-representation. The* $H_c(W)$*-submodule* I*, spanned by* U'*, and the quotient* $M_c(\tau_l)/I$ *are both irreducible.*

By claim 1, U' does not contain two-dimensional representations of W. By proposition 5.3.1, for $c > 0$ it does not contain the trivial representation either. Hence, U' is either 0 or several copies of sgn.

On the one hand, the multiplicity of sgn in U' is at most 1 (specialize to one parameter results of [Du1], p. 82 and use inequality (2)). On the other hand, according to the theorem 5.2.2, there is a singular copy of τ_l in $M_c(\mathsf{triv})$. An H_c-submodule J, generated by this copy, is not isomorphic to $M_c(\tau_l)$, since $M_c(\mathsf{triv})$ is free of rank 1 over $\mathbb{C}[\mathfrak{h}]$ meanwhile $M_c(\tau_l)$ is free of rank 2. Hence $M_c(\tau_l)$ is reducible and $U' \neq 0$. This proves that $U' \simeq \mathsf{sgn}$.

The submodule I generated by U' is isomorphic to $M_c(\mathsf{sgn})$. By theorem 5.2.1 it is irreducible.

From proposition 5.3.1 it follows that all the singular vectors in $M_c(\tau_l)/I$ are images of singular vectors in $M_c(\tau_l)$. From this fact and proposition 5.2.1 it follows, that $M_c(\tau_l)/I$ is irreducible. This completes the proof of claim 3 and of the first row in the table.

The second row in the table follows via symmetry arguments. □

Proof of theorem 3.1.1.

Claim 1. If $0 < r \equiv \pm l \pmod d$*, then* $M_c(\mathsf{triv})$ *contains a submodule isomorphic to* $L_c(\tau_l)$*, the quotient by which is irreducible.*

By theorem 5.2.2, $U' \subset M_c(\mathsf{triv})$ is isomorphic to τ_l as a representation of W, and consists of vectors of degree r. Denote an H_c-submodule of $M_c(\mathsf{triv})$ generated by U' by I_c. By definition, I_c is a module with lowest weight τ_l. By theorem 3.1.2, there are only two such modules: $M_c(\tau_l)$ and $L_c(\tau_l) \simeq M_c(\tau_l)/M_c(\mathsf{sgn})$. Since there are no singular vectors of type sgn in I_c (see theorem 5.2.2), $I_c \simeq L_c(\tau_l)$.

For eigenvalues of $\mathbf{h}$ big enough, the dimensions of the corresponding eigenspaces in $M_c(\mathsf{triv})$ and $L_c(\tau_l)$ coincide. Hence, the quotient $M_c(\mathsf{triv})/L_c(\tau_l)$ is finite dimensional. The claim follows from proposition 5.4.1.

Claim 2. If $0 < c = n + 1/2$*, then* $M_c(\mathsf{triv})$ *contains a submodule isomorphic to* $L_c(\mathsf{sgn})$*, the quotient by which is irreducible.*

By theorem 5.2.2, $U' \subset M_c(\mathsf{triv})$ is isomorphic to sgn as a W-representation, and is spanned by a vector of degree $d(2n+1)$. A submodule generated by U' is isomorphic to $M_c(\mathsf{sgn})$. By symmetry reasons and claim 1, it is irreducible, i.e. isomorphic to $L_c(\mathsf{sgn})$.

By the lowest weights reasons (proposition 5.3.1), there are no singular vectors of positive degree in $Q_c = M_c(\mathsf{triv})/M_c(\mathsf{sgn})$, and hence Q_c is irreducible. Comparing the dimensions of $\mathbf{h}$-eigenspaces one can show, that Q_c is infinite dimensional.

Claim 3. $M_c(\mathsf{triv})$ *is irreducible for all other values of* c*.*

By theorem 5.2.2, there are no singular vectors in $M_c(\mathsf{triv})$ for all other values of c. Hence, this module is irreducible. □

6.2. **Case of $d = 2m$.**

Proof of theorem 3.2.3. The proof is analogous to that of theorem 3.1.2. We will describe the action of W on the set U' of singular vectors of positive degree.

Claim 1. There are no two-dimensional irreducible subrepresentations in U'.

See proof of claim 1 of theorem 3.1.2.

Claim 2. If $(k_1, k_2) \notin E_r^+$ for all $r \equiv \pm l \pmod{2m}$ and $(k_1, k_2) \notin E_{r'}^-$ for all $r' \equiv m \pm l \pmod{2m}$, then U' does not contain any one-dimensional subrepresentations of W.

By symmetry reasons (remark 2.3.4), it is enough to prove the following fact.

Claim 2 ′. If $(k_1, k_2) \notin E_r^+$ for all $r \equiv \pm l \pmod{2m}$, then U' does not contain any W-subrepresentations isomorphic to sgn.

Suppose U' contains a W-representation isomorphic to sgn. This representation generates an H_{k_1,k_2}-submodule of $M_{k_1,k_2}(\tau_l)$, which is isomorphic to $M_{k_1,k_2}(\mathsf{sgn})$. Therefore

$$\mathrm{Hom}_{H_{k_1,k_2}}(M_{k_1,k_2}(\mathsf{sgn}), M_{k_1,k_2}(\tau_l)) \neq 0.$$

Both $M_{k_1,k_2}(\mathsf{sgn})$ and $M_{k_1,k_2}(\tau_l)$ are free over $\mathbb{C}[\mathfrak{h}]$. By theorem 5.1.6,

$$\begin{aligned} \dim \mathrm{Hom}_{H_{\mathsf{k}}}(M_{k_1,k_2}(\mathsf{sgn}), M_{k_1,k_2}(\tau_l)) \leqslant \\ \leqslant \dim \mathrm{Hom}_{\mathcal{H}}(\mathsf{KZ}(M_{k_1,k_2}(\mathsf{sgn})), \mathsf{KZ}(M_{k_1,k_2}(\tau_l))), \end{aligned} \tag{4}$$

and hence,

$$\mathrm{Hom}_{\mathcal{H}\text{-mod}}(\mathsf{KZ}(M_{k_1,k_2}(\mathsf{sgn})), \mathsf{KZ}(M_{k_1,k_2}(\tau_l))) \neq 0. \tag{5}$$

Let us compute the dimension of $\mathrm{Hom}_{\mathcal{H}\text{-mod}}(\mathsf{KZ}(M_{k_1,k_2}(\mathsf{sgn})), \mathsf{KZ}(M_{k_1,k_2}(\tau_l)))$.

Recall that in case of $W = I_2(2m)$ the Hecke algebra $\mathcal{H}(q_1, q_2)$ is spanned by two elements T_1 and T_2 with relations

$$(T_j - 1)(T_j + q_j) = 0, \quad \underbrace{T_1 T_2 \ldots T_2}_{2m \text{ multiples}} = \underbrace{T_2 T_1 \ldots T_1}_{2m \text{ multiples}},$$

In order for theorems 5.1.5 and 5.1.6 to hold, we need $q_j = e^{-2\pi i k_j}$.

The representation $\mathsf{KZ}(M_{k_1,k_2}(\tau_l))$ of $\mathcal{H}(q_1, q_2)$ corresponds to τ_l via Tits' Deformation Theorem. For $(k_1, k_2) \notin E_r^+$ for all $0 < r \equiv \pm l \pmod{2m}$, this representation is given explicitly by:

$$T_1 \mapsto \begin{pmatrix} 1 & 0 \\ c_l & -q_1 \end{pmatrix}; \quad T_2 \mapsto \begin{pmatrix} -q_2 & c_l' \\ 0 & 1 \end{pmatrix},$$

where $c_l c_l' = q_1 + q_2 + \sqrt{q_1 q_2}(\omega^l + \omega^{-l})$ and $\sqrt{q_1 q_2}$ is fixed to be $e^{-\pi i(k_1+k_2)}$ (see [GP], p. 268; note that in our the signs of the generators are different). Using the notations of [Du1], these are formulas for $-e^{2\pi i\alpha} M(\gamma_0)$ and $-e^{2\pi i\beta} M(\gamma_1)$ in the basis $\{f^{(3)}, f^{(1)}\}$.

The representation $\mathsf{KZ}(M_c(\mathsf{sgn}))$ is given explicitly by $T_1 \mapsto -q_1$ and $T_2 \mapsto -q_2$.

Hence, $\dim \mathrm{Hom}_{\mathcal{H}}(\mathsf{KZ}(M_{k_1,k_2}(\mathsf{sgn})), \mathsf{KZ}(M_{k_1,k_2}(\tau_l)) = 0$, if $(k_1, k_2) \notin E_r^+$ for all $0 < r \equiv \pm l \pmod{2m}$. This contradicts to (5), and hence U' does not contain subrepresentations isomorphic to sgn. Claim 2 is proved.

From Claims 1 and 2 it follows that $U' = 0$, when $(k_1, k_2) \notin E_r^+$ and $(k_1, k_2) \notin E_{r'}^-$ for all $r \equiv \pm l \pmod{2m}$ and all $r' \equiv m \pm l \pmod{2m}$. By proposition 5.2.1, $M_{k_1,k_2}(\tau_l)$ is irreducible for these values of (k_1, k_2). This proves part **(iv)** of the theorem.

Claim 3. If $(k_1,k_2)\in E^+_r$ for some $0<r\equiv\pm l \ (mod\, 2m)$ and $(k_1,k_2)\notin E^-_{r'}$ for all $r'\equiv m\pm l\ (mod\, 2m)$, then $U'\simeq \mathsf{sgn}$ as a representation of W. The $H_{k_1,k_2}(W)$-submodule I_{k_1,k_2}, spanned by U', and the quotient $M_{k_1,k_2}(\tau_l)/I_{k_1,k_2}$ are both irreducible.

Let us prove that U' is a direct sum of representations, isomorphic to sgn. By claim 1, U' does not contain any two-dimensional representations of W. By proposition 5.3.1, for $r>0$ it does not contain the trivial representation either. By symmetry reasons (remark 2.3.4) and claim $2'$, the W-representation U' does not contain any subrepresentations of type ε_1 and ε_2. Hence U' is either 0 or several copies of sgn.

Let us prove that the multiplicity of sgn is exactly 1. On the one hand, the multiplicity of sgn in U' is at most 1 (see inequality (4) and [Du1], p. 82). On the other hand, according to theorem 5.2.2, there is a singular copy of τ_l in $M_{k_1,k_2}(\mathsf{triv})$. An H_{k_1,k_2}-submodule J, generated by this copy, is not isomorphic to $M_{k_1,k_2}(\tau_l)$, since $M_{k_1,k_2}(\mathsf{triv})$ is free of rank 1 over $\mathbb{C}[\mathfrak{h}]$ meanwhile $M_{k_1,k_2}(\tau_l)$ is free of rank 2. Hence, $M_{k_1,k_2}(\tau_l)$ is reducible and $U'\neq 0$. This proves that $U'\simeq\mathsf{sgn}$.

The submodule I_{k_1,k_2}, generated by U', is isomorphic to $M_{k_1,k_2}(\mathsf{sgn})$. By theorem 5.2.1, it is irreducible.

Denote by Q_{k_1,k_2} the quotient $M_{k_1,k_2}(\tau_l)/I_{k_1,k_2}$. We have already proved that all the singular vectors of positive degree in Q_{k_1,k_2} are secondary, i.e. singular only "modulo I_{k_1,k_2}". By proposition 5.3.1, there are two possibilities for secondary singular vectors:

- if $(k_1,k_2)\in E^-_{r'}$ for some $r'>0$ and k_2 is negative, then Q_{k_1,k_2} might contain a singular vector of type ε_2;
- if $(k_1,k_2)\in E^-_{r'}$ for some $r'<0$ and k_1 is negative, then Q_{k_1,k_2} might contain a singular vector of type ε_1.

Suppose Q_{k_1,k_2} contains a singular vector of type ε_2. The H_{k_1,k_2}-submodule P_{k_1,k_2}, generated by this copy is isomorphic to $L_{k_1,k_2}(\varepsilon_2)\simeq M_{k_1,k_2}(\varepsilon_2)$. Denote by N the preimage of P_{k_1,k_2} in $M_{k_1,k_2}(\tau_l)$. Since P_{k_1,k_2} is generated by secondary vector, the following exact sequence does not split.

$$0\to M_{k_1,k_2}(\mathsf{sgn})\to N\to M_{k_1,k_2}(\varepsilon_2)\to 0$$

Apply KZ-functor to this sequence. The representation $\mathsf{KZ}(M_{k_1,k_2}(\mathsf{sgn}))$ is given by $T_j\mapsto -q_j$. The representation $\mathsf{KZ}(M_{k_1,k_2})(\varepsilon_1)$ is given by $T_1\mapsto -q_1$ and $T_2\mapsto 1$. To find $\mathsf{KZ}(N)$ consider the following exact sequence

$$0\to N\to M_{k_1,k_2}(\tau_l)\to J\to 0.$$

The cokernel J is supported on $\mathfrak{h}\backslash\mathfrak{h}_{reg}$, and hence $\mathsf{KZ}(J)=0$. Applying KZ-functor to the sequence above, we will get that $\mathsf{KZ}(N)\simeq\mathsf{KZ}(M_{k_1,k_2}(\tau_l))$, since the functor is exact. Hence, $\mathsf{KZ}(N)$ is given explicitly by

$$T_1\mapsto\begin{pmatrix}1 & 0\\ c_l & -q_1\end{pmatrix};\quad T_2\mapsto\begin{pmatrix}-q_2 & c'_l\\ 0 & 1\end{pmatrix}.$$

Since KZ-functor is exact, $\mathsf{KZ}(N)$ is a nontrivial extension of $\mathsf{KZ}(M_{k_1,k_2}(\varepsilon_2))$ by $\mathsf{KZ}(M_{k_1,k_2}(\mathsf{sgn}))$. It is impossible, since in every such extension $\mathrm{Tr}(T_1)=-2q_1$ meanwhile the trace of T_1 in $\mathsf{KZ}(N)$ is $1-q_1$, which is not equal to $-2q_1$ for $r'\not\equiv m\pm l$ (mod $2m$). Hence, Q_{k_1,k_2} does not contain any submodules and is irreducible.

The case of singular vector of type ε_1 is proved analogously. Claim 3 is proved.

Claim 4. Let $k_1 \geqslant k_2 > 0$ *be such that* $(k_1, k_2) \in E_r^+ \cap E_{r'}^-$ *for some* $r \equiv \pm l \ (mod\ 2m)$ *and some* $r' \equiv m \pm l \ (mod\ 2m)$*. Then* $M_{k_1,k_2}(\tau_l)$ *contains a submodule* I_{k_1,k_2}*, isomorphic to* $M_{k_1,k_2}(\mathsf{sgn})$*, and a submodule* $\widetilde{I}_{k_1,k_2}$*, isomorphic to* $M_{k_1,k_2}(\varepsilon_2)$*. If* $k_2 \in \mathbb{Z} + 1/2$*, then* $\widetilde{I}_{k_1,k_2} \supset I_{k_1,k_2}$*, otherwise* $\widetilde{I}_{k_1,k_2} \cap I_{k_1,k_2} = 0$*.*

By claim 3, $M_{k_1,k_2}(\tau_l)$ contains a submodule I_{k_1,k_2}, isomorphic to $M_{k_1,k_2}(\mathsf{sgn})$, when (k_1, k_2) is a generic point of E_r^+, and it contains a submodule $\widetilde{I}_{k_1,k_2}$, isomorphic to $M_{k_1,k_2}(\varepsilon_2)$, when (k_1, k_2) is a generic point of $E_{r'}^-$. By continuity for $(k_1, k_2) \in E_r^+ \cap E_{r'}^-$, the standard module $M_{k_1,k_2}(\tau_l)$ contains both of them.

Suppose $I_{k_1,k_2} \cap \widetilde{I}_{k_1,k_2} \neq 0$. Let $f \in I_{k_1,k_2} \cap \widetilde{I}_{k_1,k_2}$ be a vector of the minimal possible degree. Then f is a singular vector in both I_{k_1,k_2} and $\widetilde{I}_{k_1,k_2}$. The only singular vector in I_{k_1,k_2} is the one which generates this submodule. Indeed, by symmetry arguments, the structure of $M_{k_1,k_2}(\mathsf{sgn})$ is the same as the structure of $M_{-k_1,-k_2}(\mathsf{triv})$. By proposition 5.2.3, the unique singular vector contained in $M_{-k_1,-k_2}(\mathsf{triv})$ generates the whole module. Hence, if intersection of I_{k_1,k_2} and $\widetilde{I}_{k_1,k_2}$ is nonzero, then $\widetilde{I}_{k_1,k_2} \supset I_{k_1,k_2}$.

Suppose this intersection is nonzero. Then in $M_{k_1,k_2}(\varepsilon_2)$ there is a singular vector, on which $I_2(2m)$ acts by sgn. By symmetry reasons, this happens only for those (k_1, k_2), for which $M_{-k_1,k_2}(\mathsf{triv})$ contains a submodule isomorphic to $M_{-k_1,k_2}(\varepsilon_1)$. By proposition 5.2.3, this is the case if $k_2 = i + 1/2$ for some positive integer i. This proves claim 4.

Claim 5. Let (k_1, k_2) *be as in claim 4. Then the quotient* Q_{k_1,k_2} *of* $M_{k_1,k_2}(\tau_l)$ *by the submodule generated by* I_{k_1,k_2} *and* $\widetilde{I}_{k_1,k_2}$ *is irreducible.*

By proposition 5.3.1, the only possible singular vectors in Q_{k_1,k_2} are of type sgn and degree r, or of type ε_2 and degree r'.

From inequality (4) and [Du1], p. 84, it follows that the set of singular vectors in $M_{k_1,k_2}(\tau_l)$ contains only one copy of sgn and only one copy of ε_2, i.e. is isomorphic to $\tau_l \oplus \varepsilon_2 \oplus \mathsf{sgn}$ as a W-representation. Hence, all the singular vectors in Q_{k_1,k_2}, except the one, which generates this module, are secondary, i.e. singular only "modulo $\widetilde{I}_{k_1,k_2}$ and I_{k_1,k_2}".

For the values of parameters that we consider, the lowest possible degree of a vector in the ideal generated by $\widetilde{I}_{k_1,k_2}$ and I_{k_1,k_2} is r'. Hence, a vector of type ε_2 can not be secondary. Therefore, if Q_{k_1,k_2} is reducible, then it contains a singular vector of type sgn. By theorem 5.3.1, the H_{k_1,k_2}-submodule P_{k_1,k_2}, generated by this copy, is isomorphic to $L_{k_1,k_2}(\mathsf{sgn}) \simeq M_{k_1,k_2}(\mathsf{sgn})$.

For $k_2 \notin \mathbb{Z} + 1/2$ the intersection of $\widetilde{I}_{k_1,k_2}$ and I_{k_1,k_2} is zero. Suppose that Q_{k_1,k_2} has a submodule P_{k_1,k_2}. Then $Q_{k_1,k_2}/P_{k_1,k_2}$, and hence $L_{k_1,k_2}(\tau_l)$, is finite dimensional. But it is known (see [De] remark 5.7), that $L_{k_1,k_2}(\tau_l)$ is always infinite dimensional. Hence, Q_{k_1,k_2} is irreducible.

For $k_2 \in \mathbb{Z} + 1/2$ we have the following sequence of modules

$$M_{k_1,k_2}(\tau_l) \supset M_{k_1,k_2}(\varepsilon_2) \supset M_{k_1,k_2}(\mathsf{sgn}),$$

and $Q_{k_1,k_2} = M_{k_1,k_2}(\tau_l)/M_{k_1,k_2}(\varepsilon_2)$. Proof of the irreducibility of Q_{k_1,k_2} is analogous to the proof of the irreducibility of the quotient in claim 3. Claim 5 is proved.

Part **(iii)** of the theorem for $k_1 \geqslant k_2 > 0$ follows from claims 4 and 5. For all the other values of parameters the proof is analogous. □

Proof of theorem 3.2.2. **(i)** By theorem 5.2.3, the space $U' \subset M_{k_1,k_2}(\mathsf{triv})$ is isomorphic to τ_l as a representation of W and sits in degree r. The submodule I_{k_1,k_2}, spanned by U', is isomorphic to $L_{k_1,k_2}(\tau_l)$ since it does not contain any singular vectors except the one, by which it is generated. By parts **(i)** and **(iii)** of theorem 3.2.3, $L_{k_1,k_2}(\tau_l) \simeq M_{k_1,k_2}(\tau_l)/M_{k_1,k_2}(\mathsf{sgn})$.

For eigenvalues of $\mathbf{h}$ big enough, the dimensions of the corresponding eigenspaces in $M_{k_1,k_2}(\mathsf{triv})$ and $L_{k_1,k_2}(\tau_l)$ coincide. Hence, the quotient $M_{k_1,k_2}(\mathsf{triv})/L_{k_1,k_2}(\tau_l)$ is finite dimensional. The statement follows from proposition 5.4.1.

(ii) By theorem 5.2.3, the space $U' \subset M_{k_1,k_2}(\mathsf{triv})$ is isomorphic to ε_2 as a W-representation. It is easy to see that it generates a submodule isomorphic to $M_{k_1,k_2}(\varepsilon_2)$.

Consider the quotient $Q_{k_1,k_2} = M_{k_1,k_2}(\mathsf{triv})/M_{k_1,k_2}(\varepsilon_2)$. From the proposition 5.3.1 it follows that Q_{k_1,k_2} does not contain any singular vectors and hence is irreducible. Comparing dimensions of eigenspaces of $\mathbf{h}$ in $M_{k_1,k_2}(\varepsilon_2)$ and in $M_{k_1,k_2}(\mathsf{triv})$, one can show that Q_{k_1,k_2} is infinite dimensional.

(iii) Analogous to part **(ii)**.

(iv) Consider the case of $(k_1, k_2) \in E_r^+ \cap L_i^1$ with $k_1 < k_2$. The other case is proved analogously.

By theorem 5.2.3, the space $U' \subset M_{k_1,k_2}(\mathsf{triv})$ is isomorphic to $\tau_l \oplus \varepsilon_2$ as a W-representation.

A submodule I_{k_1,k_2} of $M_{k_1,k_2}(\mathsf{triv})$, generated by a singular vector of type ε_2, is isomorphic to $M_{k_1,k_2}(\varepsilon_2)$. By symmetry reasons, $M_{k_1,k_2}(\varepsilon_2)$ has the same structure as $M_{-k_1,k_2}(\mathsf{triv})$ and hence, by part **(i)**, it contains a singular copy of τ_l, which generates a submodule isomorphic to $L_{k_1,k_2}(\tau_l)$. So we have the following picture:

$$M_{k_1,k_2}(\mathsf{triv}) \supset M_{k_1,k_2}(\varepsilon_2) \supset L_{k_1,k_2}(\tau_l).$$

By symmetry arguments, since $M_{-k_1,k_2}(\mathsf{triv})/L_{-k_1,k_2}(\tau_{m-l})$ is irreducible, so is the quotient $M_{k_1,k_2}(\varepsilon_2)/L_{k_1,k_2}(\tau_l)$.

Consider the quotient $Q_{k_1,k_2} = M_{k_1,k_2}(\mathsf{triv})/M_{k_1,k_2}(\varepsilon_2)$. Comparing the dimensions of $\mathbf{h}$-eigenspaces in $M_{k_1,k_2}(\mathsf{triv})$ and $M_{k_1,k_2}(\varepsilon_2)$, one gets that Q_{k_1,k_2} is infinite dimensional.

Since Q_{k_1,k_2} has the lowest weight triv, its irreducible quotient is isomorphic to $L_{k_1,k_2}(\mathsf{triv})$. For all (k_1, k_2) the module $L_{k_1,k_2}(\mathsf{triv})$ is defined as a quotient of $M_{k_1,k_2}(\mathsf{triv})$ by the kernel of Shapovalov form. By part **(i)**, for generic $(k_1, k_2) \in E_r^+$ the irreducible quotient has dimension r^2 and hence $\dim L_{k_1,k_2}(\mathsf{triv}) \leqslant r^2$.

Hence Q_{k_1,k_2} is reducible and contains singular vectors. Note that these vectors are secondary, i.e. singular only "modulo $M_{k_1,k_2}(\varepsilon_2)$". By the table of lowest $\mathbf{h}$-weights (proposition 5.3.1), there exit the following possibilities for singular vectors: vectors of type τ_l sitting in degree r, vectors of type ε_1 sitting in degree $2mk_2$ or vectors of type sgn in degree $2r$.

The sequence of the dimensions of $\mathbf{h}$-eigenspaces in Q_{k_1,k_2} is

$$(1, 2, 3, \ldots, 2k_1m, 2k_1m, 2k_1m, \ldots). \tag{6}$$

Since $L_{k_1,k_2}(\mathsf{triv})$ is a finite dimensional $\mathfrak{sl}_2$-module, the sequence of dimensions of its $\mathbf{h}$-eigenspaces is finite and symmetric with respect to the 0-eigenspace. From the table of lowest $\mathbf{h}$-weights (proposition 5.3.1) it follows that $\mathbf{h}$ acts by 0 on polynomials of degree $r - 1$. In order to make the sequence (6) symmetric, Q_{k_1,k_2} has to contain a unique up to proportionality singular vector, which is of type ε_1 and sits in degree $2mk_2$. This vector generates a submodule P_{k_1,k_2} isomorphic to

$L_{k_1,k_2}(\varepsilon_1)$. From the sequence of dimensions of $\mathbf{h}$-eigenspaces in $Q_{k_1,k_2}/P_{k_1,k_2}$ it follows that this quotient is is irreducible.

(v) Consider the case $(k_1, k_2) \in E_r^+ \cap L_i^1$. By theorem 5.2.3, there exists a singular vector of type ε_2 in $M_{k_1,k_2}(\mathsf{triv})$. It generates a submodule isomorphic to $M_{k_1,k_2}(\varepsilon_2)$. This submodule is irreducible since, by symmetry arguments, it has the same structure as $M_{-k_1,k_2}(\mathsf{triv})$.

Consider the quotient $Q_{k_1,k_2} = M_{k_1,k_2}(\mathsf{triv})/M_{k_1,k_2}(\varepsilon_2)$. By theorem 5.2.3, there is a singular copy of τ_l in Q_{k_1,k_2}. Denote by P_{k_1,k_2} the submodule generated by this copy. By definition, this is a module of lowest weight τ_l. By theorem 3.2.3, for these values of parameters there are three such modules: $M_{k_1,k_2}(\tau_l)$, $M_{k_1,k_2}(\tau_l)/M_{k_1,k_2}(\mathsf{sgn})$ and $L_{k_1,k_2}(\mathsf{triv}) \simeq M_{k_1,k_2}(\mathsf{triv})/M_{k_1,k_2}(\varepsilon_2)$. Since there are no singular vectors of type sgn and ε_2 in Q_{k_1,k_2}, the submodule P_{k_1,k_2} is isomorphic to $L_{k_1,k_2}(\tau_l)$.

Using the symmetry of the sequence of the dimensions of $\mathbf{h}$-eigenspaces, one can show that $Q_{k_1,k_2}/P_{k_1,k_2}$ is irreducible.

(vi) By theorem 5.2.3, in $M_{k_1,k_2}(\mathsf{triv})$ there are singular vectors of types ε_1 and ε_2. They generates submodules isomorphic to $M_{k_1,k_2}(\varepsilon_1)$ and $M_{k_1,k_2}(\varepsilon_2)$ respectively. These two modules intersect each other by a submodule isomorphic to $M_{k_1,k_2}(\mathsf{sgn})$ and generated by a singular vector. By symmetry reasons, the quotients $M_{k_1,k_2}(\varepsilon_1)/M_{k_1,k_2}(\mathsf{sgn})$ and $M_{k_1,k_2}(\varepsilon_2)/M_{k_1,k_2}(\mathsf{sgn})$ are irreducible. The quotient of $M_{k_1,k_2}(\mathsf{triv})$ by the union of $M_{k_1,k_2}(\varepsilon_1)$ and $M_{k_1,k_2}(\varepsilon_2)$ is irreducible for the same reasons as in parts **(iv)** and **(v)**.

(viii) By theorem 5.2.3, the space U' is empty and hence, by proposition 5.2.1 $M_{k_1,k_2}(\mathsf{triv})$ is irreducible. $\square$

Acknowledgements. I am very grateful to P. Etingof for suggesting me this problem and for many useful conversations. I would also like to thank I. Scherbak and S. Chmutov for many comments.

References

[BEG] Yu. Berest, P. Etingof, V. Ginzburg, *Finite dimensional representations of rational Cherednik algebras,* Preprint arXiv:math.RT/0208138.

[BEG1] Yu. Berest, P. Etingof, V. Ginzburg, *Cherednik algebras and differential operators on quasi-invariants,* Preprint arXiv:math.QA/0111005.

[BMR] M. Broué, G. Malle, R. Rouquier, *Complex reflection groups, braid groups,Hecke algebras,* J. reine angew. Math 500(1998), 127-190.

[Br] E. Brieskorn, *Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen komplexen Spiegelungsgruppe*, Invent. Math. 12(1971), 57-61.

[CE] T. Chmutova, P. Etingof, *On some representations of the rational Cherednik algebra,* Preprint arXiv:math.RT/0303194

[De] C. Dezélée, *Représentations de dimension finie de l'algèbre de Cherednik rationnelle*, Preprint arXiv:math.RT/0111210.

[DJO] C. F. Dunkl, M. F. E. De Jeu, E. M. Opdam, *Singular polynomials for finite reflection groups.*, Trans. Amer. Math. Soc.346 (1994) 237-256.

[DO] C. F. Dunkl, E. M. Opdam, *Dunkl operators for complex reflection groups*, Preprint math.RT/0108185.

[Du] C. F. Dunkl, *Differential-difference operators associated to reflection groups.* Trans. Amer. Math. Soc. 311(1989) 167-183.

[Du1] C. F. Dunkl, *Monodromy of hypergeometric functions for dihedral groups,* Integral Transform. Spec. Funct. 1 (1993), no. 2, 75-86.

[EG] P. Etingof, V. Ginzburg, *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism,* Invent. Math. 147(2002), 243-348, Preprint arXiv:math.AG/0011114.

[Go] I. Gordon, *On the quotient ring by diagonal harmonics,* arXiv:math.RT/0208126.

[GP] M. Geck, G. Pfeiffer, *Characters of finite Coxeter groups and Iwahori-Hecke Algebras*, LMS Monographs 21, (2000).

[GGOR] V. Ginzburg, N. Guay, E. Opdam, R. Rouquier, *On category $\mathcal{O}$ for rational Cherednik algebras*, Preprint math.RT/0212036.

Department of Mathematics, Harvard University, Cambridge, MA 02138, USA
E-mail address: `chmutova@math.harvard.edu`